

# HOME FALLS AMONG PEOPLE 65 YEARS AND OLDER: CIRCUMSTANCES AND PATTERNS OF FALLS:

*RESULTS FROM THE CHUPADOM SURVEY - 2018*



Marion Torres, Gaëlle Pedrono, Linda Lasbeur, Laure Carcaillon-Bentata, Annabel Rigou, Nathalie Beltzer

(Santé publique France, Saint Maurice, France)

## **INTRODUCTION**

- Falls are responsible for important morbidity and mortality → major public health issue in elderly people
- In France, in 2010:
  - 22% of people aged between 55-85 fall at least once
  - Falls accounted for 85% of emergencies for Home and Leisure Injuries (HLIs) among people aged 65 and over
- 7/10 falls occurred at home
- Currently, few data exist to characterize the heterogeneity of the circumstances of falls which would be helpful to set up targeted prevention actions

## **OBJECTIVES**

- To describe the characteristics and circumstances surrounding falls occurring at home among people aged 65 and over hospitalized, and to identify profiles of fallers


## METHODS

- ChuPADom: prospective multicenter survey conducted in 2018 in 7 volunteer French hospitals
- For each, all persons aged 65 and over hospitalized within a 6-month period following a fall at home interviewed
- Data collected during a face-to-face interview at the hospital by a referring hospital investigator
- Following a description of the sample, patterns of falls were identified by Multiple Correspondence of Analysis (MCA) and Ascending Hierarchical Classification (CAH)




## RESULTS

- 1 467 patients included (69% female, mean age 84,5y)




- Morning (6am-1pm)
- Afternoon (1pm-6pm)
- Evening (6pm-11pm)
- Night (11pm-6am)

Time of the fall



History of falls and risk-taking



Circumstances of the fall

## RESULTS: Profiles of fallers

### Profile 1 (n=83, 5,7%):

Falls: high, moderate to intense, outdoor or garage, risk taking


- Men, 65-74 yrs,
- Autonomous,
- Physical activity,
- homeowner, house

### Profile 2 (n=67, 4,6%):

- Falls: stairs, moderate to intense,
- pre-fragile,
- Autonomous,
- Home,
- Married/Cohabiting,
- Education or more,
- 65-84 years old

### Profile 5 (n=49, 3,0%):

- Missing data for housing adapted to health status, perceived health, autonomy, circumstances of the fall...
- Retirement home,
- 90 yrs and older



### Factors used to build profiles

### Profile 3 (n=476, 32,4%):

- Autonomous,
- Fall: outdoor, moderate to intense, loss of balance or low height,
- 65-84 years,
- Physical activity, -very good perceived health,
- Homeowner,
- Housing adapted to health status

### Profile 4 (n=792, 54,0%):

- Dependent,
- Motor impairment,
- No physical activity,
- Retirement home,
- Fall: bedroom and living room, rest/transfer, mecha NP, night,
- Widowed, Women, 85+ years old,
- Poor perceived health, neurological diseases, CVD, psychological diseases, fatigue

## DISCUSSION/CONCLUSION

- These results highlight the heterogeneity of the circumstances in which elderly people fall
- Profiles used for the anti-fall plan launched in 2022 by the French government to implement prevention actions
- Results of this study will be enriched by the analysis of the second part of this survey with consequences of the fall one year after



ELSEVIER

Contents lists available at ScienceDirect

Injury

journal homepage: [www.elsevier.com/locate/injury](http://www.elsevier.com/locate/injury)



Identifying profiles of people aged 65 and over who fall at home and associated falling-induced injuries: The French ChuPADom study

Marion J Torres\*, Gaëlle Pédrone, Annabel Rigou, Laure Carcaillon-Bentata, Nathalie Beltzer

Direction des Maladies Non Transmissibles et Traumatismes, Santé Publique France, Unité Pathologies, Populations et Traumatismes, French National Public Health Agency, 12 rue du Val d'Osne, Saint-Maurice Cedex 94415, France